

**Supplemental Table S1**

<b>Ineffective compounds</b>	<b>Tested on</b>	<b>Concentration range used</b>
<b>Doxycycline</b>	<b>BJ</b>	<b>25 – 200 <math>\mu</math>M</b>
<b>Diphenyleneiodonium chloride</b>	<b>BJ</b>	<b>0.5 – 10 <math>\mu</math>M</b>
<b>Melatonin</b>	<b>BJ</b>	<b>200 – 400 <math>\mu</math>M</b>
<b>Aspartame</b>	<b>BJ</b>	<b>200 – 400 <math>\mu</math>M</b>
<b>Glucosamine</b>	<b>BJ</b>	<b>10 – 15 mM</b>
<b>Quercetin</b>	<b>BJ</b>	<b>10 – 100 <math>\mu</math>M</b>
<b>Dasatinib</b>	<b>BJ</b>	<b>0.1 – 1 <math>\mu</math>M</b>
<b>Chloroquine</b>	<b>BJ, MRC-5</b>	<b>100 – 200 <math>\mu</math>M</b>
<b>Erythromycin</b>	<b>BJ, MRC-5</b>	<b>100 – 200 <math>\mu</math>M</b>
<b>Clarithromycin</b>	<b>BJ, MRC-5</b>	<b>100 – 200 <math>\mu</math>M</b>
<b>Rapamycin</b>	<b>MRC-5</b>	<b>50 – 500 nM</b>
<b>Lycopene</b>	<b>MRC-5</b>	<b>25 – 50 <math>\mu</math>M</b>
<b>Alpha-lipoic acid</b>	<b>MRC-5</b>	<b>25 – 50 <math>\mu</math>M</b>