

SUPPLEMENTARY TABLES

Supplementary Table 1. The primers for real-time QPCR.

Gene	Forward or Reverse	Primer sequence
PVT1	Forward	5'- GCCCCTTCTATGGGAATCACTA -3'
	Reverse	5'- GGGGCA GAGATGAAATCGTAAT -3'
BCLAF1	Forward	5'- ATGAGACGACCTTATGGGTACA -3'
	Reverse	5'- AGAGTGCCTTCTATTCCA GACAG -3'
miR-194-5p	Forward	5'- GCCGCTGTAACA GCAACTCCAT -3'
	Reverse	5'- GTGCAGGGTCCGAGGT -3'
GAPDH	Forward	5'- CGCTCTCTGCTCCTCCTGTTC-3'
	Reverse	5'-ATCCGTTGACTCCGACCTTCA C-3'
U6	Forward	5'- CTCGCTTCGGCA GCACA -3'
	Reverse	5'- ACGCTTCACGAATTTGCGT-3

BCLAF1: BCL2 associated transcription factor 1.

Supplementary Table 2. Antibodies used for western blots.

Antibody	Description	Dilution	Supplier	Country
Anti-CDC20	Mouse monoclonal	1:1000	Santa Cruz	USA
Anti- CD44	Mouse monoclonal	1:1000	Santa Cruz	USA
Anti-Met	Mouse monoclonal	1:1000	Santa Cruz	USA
Anti-β-catenin	Mouse monoclonal	1: 1000	Santa Cruz	USA
Anti-Cyclin D1	Rabbit monoclonal	1: 1000	Beyotime	China
Anti-BCLAF1	Rabbit polyclonal	1: 500	BBI	China
Anti-BCLAF1	Mouse monoclonal	1: 500	Santa Cruz	USA
Anti-β-Actin	Mouse monoclonal	1: 1000	Beyotime	China
Anti-BCLAF1	Rabbit Polyclonal	1: 500	Proteintech	USA

CDC20: cell division cycle 20; BCLAF1: BCL2 associated transcription factor 1; CD44: CD44 molecule; Met: MET proto-oncogene, receptor tyrosine kinase; β-catenin: catenin beta 1.